

62719-175

12/17/2012

1/30



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

Nneka Breaux, Ph.D.  
Dow AgroSciences LLC  
9330 Zionsville Road  
Indianapolis, IN 46268

DEC 17 2012

Subject: Product Name: Snapshot 2.5 TG  
Proposed Alternate Brand Name: Snapshot DG Specialty Herbicide  
EPA Reg. No. 62719-175  
Label Notification per PRN 98-10

Dear Dr. Breaux,

The Agency is in receipt of your application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for Snapshot 2.5 TG (EPA Reg. No. 62719-175) dated June 18, 2012. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please contact Emily Hartman of my staff at (703) 347-0189 or [hartman.emily@epa.gov](mailto:hartman.emily@epa.gov).

Sincerely,

A handwritten signature in black ink that reads "Kathryn V. Montague".

Kathryn V. Montague, Project Manager 23  
Herbicide Division  
Registration Division  
Office of Pesticide Programs



United States  
Environmental Protection Agency  
Washington, DC 20460

 Registration  
 Amendment  
 Other

OPP Identifier Number

### Application for Pesticide - Section I

1. Company/Product Number Dow AgroSciences/62719-175	2. EPA Product Manager Katherine Montague	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Dow AgroSciences/Snapshot® 2.5 TG	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: <b>NOTIFICATION</b> EPA Reg. No. _____ Product Name _____ <b>DEC 17 2012</b>	

### Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

**Explanation:** Use additional page(s) if necessary. (For section I and Section II.)

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

### Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal	<input type="checkbox"/> Plastic
				<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
* Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt	No. per container
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/>	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

### Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Nneka T. Breaux, Ph.D. (nbreaux@dow.com)	Title Regulatory Manager	Telephone No. (include Area Code) (317) 337-3248
2. Signature 		3. Title Regulatory Manager
4. Typed Name Nneka T. Breaux, Ph.D. @Trademark of Dow AgroSciences LLC		5. Date June 18, 2012
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)





### User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

### First Aid

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for treatment advice.

**If swallowed:** Call a Poison Control Center or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a Poison Control Center or doctor. Do not give anything by mouth to an unconscious person.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a Poison Control Center or doctor for further treatment advice.

Have the product container or label with you when calling a Poison Control Center or doctor or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

### Environmental Hazards

This pesticide is extremely toxic to freshwater marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply in a manner which will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

### (Storage and Disposal for rigid containers 5 gal or less)

#### Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide Storage:** Store in original container only. In case of spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

**(Storage and Disposal for nonrigid containers any size)****Storage and Disposal**

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide Storage:** Store in original container only. In case of spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available, or dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

**(Storage and Disposal for refillable rigid containers larger than 5 gal)****Storage and Disposal**

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide Storage:** Store in original container only. In case of spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Handling:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

**(Storage and Disposal for nonrefillable rigid containers larger than 5 gal)****Storage and Disposal**

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide Storage:** Store in original container only. In case of spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Refer to label booklet for Directions for Use.



8/30

(Label booklet cover):

# Snapshot<sup>®</sup> 2.5 TG

## Specialty Herbicide

**A selective preemergence herbicide for control of certain broadleaf weeds and annual grasses in:**

- Landscape Ornamentals
- Christmas Tree Plantations
- Container Grown Ornamentals
- Field Grown Ornamentals
- Groundcovers/Perennials
- Non-Bearing Fruit and Nut Trees
- Non-Bearing Vineyards
- Non-Cropland

**Active Ingredients:**

trifluralin: $\alpha, \alpha, \alpha$ -trifluoro-2,6-dinitro- <i>N</i> , N-dipropyl- <i>p</i> -toluidine.....	2.0%
isoxaben: <i>N</i> -[3-(1-ethyl-1-methylpropyl)-5- isoxazolyl]-2,6-dimethoxybenzamide and isomers.....	0.5%
Other Ingredients.....	97.5%
Total.....	100.0%

Contains 1.25 lb active ingredient per 50 lb bag.

### Keep Out of Reach of Children

## CAUTION PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to inside of label booklet for additional precautionary information including Directions for Use.

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at [www.dowagro.com](http://www.dowagro.com).


Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-175

EPA Est. \_\_\_\_\_

®Trademark of Dow AgroSciences LLC

Produced for  
Dow AgroSciences LLC





(Page 1 through end):

**Precautionary Statements****Hazards to Humans and Domestic Animals****CAUTION**

**Causes Moderate Eye Irritation • Harmful If Swallowed Or Inhaled • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reaction In Some Individuals**

**Avoid breathing dust or spray mist and contact with eyes or clothing. Wash thoroughly with soap and water after handling.**

**Personal Protective Equipment (PPE)**

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**Engineering Controls**

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**User Safety Recommendations**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

**First Aid**

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for treatment advice.

**If swallowed:** Call a Poison Control Center or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a Poison Control Center or doctor. Do not give anything by mouth to an unconscious person.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a Poison Control Center or doctor for further treatment advice.

Have the product container or label with you when calling a Poison Control Center or doctor or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

**Environmental Hazards**

This pesticide is extremely toxic to freshwater marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply in a manner which will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

11/30

## Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements of this label about personal protective equipment (PPE), and restricted entry interval. The requirements in the box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

### Non-Agricultural Use Requirements

The requirements of this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

For applications in landscape settings and in non-cropland sites, do not enter or allow others to enter the treated area until dusts have settled.

### Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Store in original container only. In case of spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

#### Nonrefillable rigid containers 5 gallons or less:

**Container Handling:** Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.



days of application. If rainfall or irrigation does not occur within 3 days of application and tillage is possible, Snapshot 2.5 TG may be activated using cultivation equipment capable of uniformly mixing the herbicide into the upper 1 to 2 inches of soil. Failure to activate Snapshot 2.5 TG within 3 days of application may result in erratic control of annual grasses. Do not apply when wind conditions favor drift of Snapshot 2.5 TG granules from the target area.

Repeat applications at 150 lb per acre and higher should not be made sooner than 60 days after a previous application of Snapshot 2.5 TG. Do not apply over 600 pounds per acre total of Snapshot 2.5 TG within a 12-month period.

**Tolerance of Turfgrass Adjacent to Ornamental Plantings:** Accidental application to turf may occur when applying Snapshot 2.5 TG to ornamental plantings. Snapshot 2.5TG is not recommended for weed control in turfgrasses, but turfgrasses are generally tolerant to small amounts of this product that fall outside of the intended area of application.

Do not aerially apply Snapshot 2.5TG.

**Treatment Species Not Listed on the Label for Snapshot 2.5 TG**

Users who wish to use Snapshot 2.5 TG on plant species not recommended on this label may determine the suitability for use by treating a small number of such plants at a recommended rate. Prior to treatment of larger areas, the treated plants should be observed for any sign of herbicidal injury for during 30 to 60 days of normal growing conditions to determine if the treatment is safe to the target plant species. The user assumes responsibility for any plant damage or other liability resulting from use of Snapshot 2.5 TG on plant species not recommended on this label.

**Application Instructions**

Apply Snapshot 2.5 TG using a drop or rotary-type spreader designed to apply granular herbicides or insecticides. Calibrate application equipment prior to use according to manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly. Do not use spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or crop injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at right angles to the first.


**Application Techniques for Applying Snapshot 2.5 TG**

- When using a drop-type spreader, a splashboard mounted under the hopper will provide more even granule distribution.
- A chain fastened to the side of the spreader and allowed to drag on the soil surface can be used to mark the edge of the treated swath and help prevent skips or overlaps.
- For treating smaller areas or rows of nursery stock or ornamental beds, a hand held or push-type rotary applicator such as a whirlybird or cyclone unit is recommended. For hand held units, walk and turn the crank at a constant rate of speed.
- A shaker-type applicator made from a small container with holes punched in the bottom is recommended for small, difficult to treat areas. Carefully measure the amount of product needed to avoid over application.
- The weight-to-volume conversion table below provides equivalent amounts of Snapshot 2.5 TG.

**Broadcast Rates**

Rate lb/Acre	Amount/1000 sq ft		Amount/100 sq ft	
	(lb)	(quarts)	(lb)	(cups)
100	2.30	1.75	0.23	0.7
150	3.45	2.60	0.35	1.0
200	4.60	3.50	0.46	1.4

1 lb of Snapshot 2.5 TG = 0.75 quart = 3 cups



14/30

**Spreader Settings as a Guide for Calibration**

**Note:** These settings are provided as a suggested starting point in calibrating each individual spreader. Since no two spreaders are alike, these settings are not intended to be used as absolute recommendations by Dow AgroSciences or the spreader manufacturer.


Settings for Warren T-7 II Spreader			
Target Rate of Snapshot 2.5 TG			
Speed (mph)	100 lb/acre	150 lb/acre	200 lb/acre
2.0	2 1/2	3 1/4	3 3/4
2.5	3	3 1/2	4 1/4
3.0	3 1/4	4	5

Warren spreader settings are displayed in half-number increments only. The suggested settings that indicate quarter settings can be accomplished by placing the calibration arm between the half number marks on the spreader.

**Weeds Controlled or Suppressed**

Weeds controlled when applied at 100 lb per acre (2.3 lb per 1000 sq ft):

Common Name	Scientific Name
aster, slender	<i>Aster exilis</i>
barnyardgrass	<i>Echinochloa crus-galli</i>
bluegrass, annual	<i>Poa annua</i>
bursage, annual	<i>Ambrosia acanthicarpa</i>
celery, wild	<i>Apium leptophyllum</i>
chickweed, common	<i>Stellaria media</i>
clover, white	<i>Trifolium repens</i>
crabgrass	<i>Digitaria</i> spp.
cudweed, purple	<i>Gnaphalium purpureum</i>
cupgrass, southwestern	<i>Eriochloa gracilis</i>
fiddleneck, coast	<i>Amsinckia intermedia</i>
filaree, redstem	<i>Erodium cicutarium</i>
fleabane, blackleaved	<i>Conyza bonariensis</i>
fleabane, dwarf	<i>Conyza ramosissima</i>
foxtail, yellow	<i>Setaria glauca</i>
groundcherry, lanceleaf	<i>Physalis lanceifolia</i>
henbit	<i>Lamium amplexicaule</i>
horseweed	<i>Conyza canadensis</i>
junglerice	<i>Echinochloa colonum</i>
knotweed, prostrate	<i>Polygonum aviculare</i>
lambsquarters, common	<i>Chenopodium album</i>
mallow, little	<i>Malva parviflora</i>
mustard, Indian	<i>Brassica juncea</i>
mustard, wild	<i>Sinapis arvensis</i>
nightshade, black	<i>Solanum nigrum</i>
oat, wild	<i>Avena fatua</i>
panicum, fall	<i>Panicum dichotomiflorum</i>
pepperweed, Virginia	<i>Lepidium virginicum</i>
pigweed	<i>Amaranthus</i> spp.
pineappleweed	<i>Matricaria matricarioides</i>
plantain, slender	<i>Plantago elongata</i>
purslane, common	<i>Portulaca oleracea</i>
radish, wild	<i>Raphanus raphanistrum</i>
ragweed, common	<i>Ambrosia artemisiifolia</i>



rocket, London	<i>Sisymbrium irio</i>
rockpurslane, desert	<i>Calandrinia ciliata</i>
shepherdspurse	<i>Capsella bursa-pastoris</i>
sibara	<i>Sibara virginica</i>
smartweed, Pennsylvania	<i>Polygonum pennsylvanicum</i>
sowthistle, annual	<i>Sonchus oleraceus</i>
speedwell, purslane	<i>Veronica peregrina</i>
sprangletop, Mexican	<i>Leptochloa univervia</i>
telegraphplant	<i>Heterotheca grandiflora</i>
thistle, Russian	<i>Salsola iberica</i>
witchgrass	<i>Panicum capillare</i>

**Weeds controlled when applied at 150 lb per acre (3.5 lb per 1000 sq ft):**

<b>Common Name</b>	<b>Scientific Name</b>
aster, heath	<i>Aster ericoides</i>
bittercress	<i>Cardamine oligosperma</i>
bittercress, hairy	<i>Cardamine hirsuta</i>
brassbuttons, southern	<i>Cotula australis</i>
carrot, wild	<i>Daucus carota</i>
chamberbitter	<i>Phyllanthus urinaria</i>
chickweed, mouseear	<i>Cerastium vulgatum</i>
dandelion	<i>Taraxacum officinale</i>
eclipta	<i>Eclipta prostrata</i>
fireweed	<i>Erechtites hieracifolia</i>
foxtail, giant	<i>Setaria faberi</i>
galinsoga, hairy	<i>Galinsoga ciliata</i>
geranium, Carolina	<i>Geranium carolinianum</i>
groundsel, common	<i>Senecio vulgaris</i>
ladysthumb	<i>Polygonum persicaria</i>
lettuce, prickly	<i>Lactuca serriola</i>
lovegrass	<i>Eragrostis</i> spp.
mallow, dwarf	<i>Malva rotundifolia</i>
marestalk	<i>Hippuris vulgaris</i>
mayweed	<i>Anthemis cotula</i>
morningglory, ivyleaf	<i>Ipomoea hederacea</i>
mustard, black	<i>Brassica nigra</i>
pennywort	<i>Hydrocotyle</i> spp.
phyllanthus, long-stalk	<i>Phyllanthus tenellus</i>
plantain, bracted	<i>Plantago aristata</i>
plantain, broadleaf	<i>Plantago major</i>
plantain, buckhorn	<i>Plantago lanceolata</i>
pokeweed, common	<i>Phytolacca americana</i>
rockpurslane, redmaids	<i>Calandrinia ciliata</i> var. <i>menziesii</i>
ryegrass, annual	<i>Lolium multiflorum</i>
sida, prickly	<i>Sida spinosa</i>
sorrell, red	<i>Rumex acetosella</i>
speedwell, thymeleaf	<i>Veronica serpyllifolia</i>
spurge, hyssop	<i>Euphorbia hyssopifolia</i>
spurge, spotted	<i>Euphorbia maculata</i>
sweetclover, yellow	<i>Melilotus officinalis</i>
tansymustard, green	<i>Descurainia pinnata</i>
velvetleaf	<i>Abutilon theophrasti</i>
woodsorrel, yellow	<i>Oxalis stricta</i>

**Weeds controlled when applied at 200 lb per acre (4.6 lb per 1000 sq ft):**

<b>Common Name</b>	<b>Scientific Name</b>
barley, hare	<i>Hordeum leporinum</i>
bromegrass	<i>Bromus</i> spp.
burclover, California	<i>Medicago polymorpha</i>
cheat	<i>Bromus secalinus</i>
datura	<i>Datura</i> spp.
dogfennel	<i>Eupatorium capillifolium</i>
eveningprimrose	<i>Oenothera</i> spp.
fescue, rattail	<i>Vulpia myuros</i>
filaree, whitestem	<i>Erodium moschatum</i>
goosefoot, nettleleaf	<i>Chenopodium murale</i>
goosegrass	<i>Eleusine indica</i>
jimsonweed	<i>Datura stramonium</i>
knotweed, silversheath	<i>Polygonum argyrocoleon</i>
kochia	<i>Kochia scoparia</i>
medic, black	<i>Medicago lupulina</i>
mullein, turkey	<i>Eremocarpus setigerus</i>
nettle, burning	<i>Urtica urens</i>
nettle, stinging	<i>Urtica dioica</i>
oxtongue, bristly	<i>Picris echioides</i>
pimpernel, scarlet	<i>Anagallis arvensis</i>
sandbur, field	<i>Cenchrus incertus</i>
signalgrass	<i>Brachiaria</i> spp.
sowthistle, spiny	<i>Sonchus asper</i>
spurge, petty	<i>Euphorbia peplus</i>
spurge, prostrate	<i>Euphorbia humistrata</i>
stinkgrass	<i>Eragrostis cilianensis</i>
sunflower	<i>Helianthus</i> spp.
swinecress	<i>Coronopus didymus</i>
thistle, musk	<i>Carduus nutans</i>
willoweed, panicle	<i>Epilobium paniculatum</i>
woodsorrel, creeping	<i>Oxalis corniculata</i>

**Weeds partially controlled or suppressed when applied at 200 lb per acre (4.6 lb per 1000 sq ft):**

<b>Common Name</b>	<b>Scientific Name</b>
bindweed, field	<i>Convolvulus arvensis</i>
carpetweed	<i>Mollugo verticillata</i>
dock, curly	<i>Rumex crispus</i>
johnsongrass (seedling)	<i>Sorghum halepense</i>
mallow, Venice	<i>Hibiscus trionum</i>
milkweed, honeyvine	<i>Ampelamus albidus</i>
morningglory, tall	<i>Ipomoea purpurea</i>
panicum, Texas	<i>Panicum texanum</i>
pusley, Florida	<i>Richardia scabra</i>
shattercane	<i>Sorghum bicolor</i>

**Uses**

**Ornamental Plantings**

Snapshot 2.5 TG is recommended as a preemergence treatment for control of certain broadleaf weeds and annual grasses in container grown ornamentals, landscape ornamentals, field grown ornamentals, ground covers/perennials, Christmas trees, non-bearing fruit and nut crops, and non-bearing vineyards. Apply Snapshot 2.5 TG anytime prior to germination of target weeds, or immediately after cultivation.

17/30

Refer to the General Information section of this label for General Use Precautions and information on application, application rates and weeds controlled.

**Special Use Precautions:**

To avoid possible injury, do not apply Snapshot 2.5 TG to:

- nursery, forest, or Christmas tree: seedling beds, cutting beds, or transplant beds.
- nursery seedbeds or forest or Christmas tree seedling transplant beds.
- unrooted liners or cuttings that have been planted in pots for the first time.
- pots less than four inches wide.
- bedding plants or areas where bedding plants will be planted or transplanted within one year following application.
- groundcovers until they are established and well rooted.

Do not apply Snapshot 2.5 TG to newly transplanted ornamentals, nursery stock, Christmas trees, groundcovers, non-bearing fruit and nut trees and non-bearing vineyards until soil or potting media has been settled by packing and irrigation or rainfall and no cracks are present or injury may occur.


Repeat applications at 150 lb per acre and higher should not be made sooner than 60 days after a previous application of Snapshot 2.5 TG. Do not apply over 600 lb per acre total of Snapshot 2.5 TG within a 12-month period.

Where loss of stand has occurred in field grown ornamentals, use untreated soil as fill around roots when replacing plants or injury may occur.

**Note:** Injury on the following plant species has been observed following applications of Snapshot 2.5 TG and use is not recommended.

<b>Common Name</b>	<b>Scientific Name</b>
ballerina cranesbill	<i>Geranium cinereum 'Ballerina'</i>
bertram anderson lingwort	<i>Pulmonaria longifolia</i>
bugle	<i>Ajuga</i> spp.
cajeput tree	<i>Melaleuca quinquenervia</i>
candytuft	<i>Iberis</i> spp.
Carolina rhododendron	<i>Rhododendron carolinianum</i>
dwarf burning bush	<i>Euonymus alatus 'compacta'</i>
foxglove	<i>Digitalis purpurea</i>
green yucca	<i>Yucca recurvifolia</i>
hydrangea	<i>Hydrangea</i> spp.
luxuriant bleeding heart	<i>Dicentra luxuriant</i>
mountain sandwort	<i>Arenaria montana</i>
mustard	<i>Brassica</i> spp.
oak leaf acanthus	<i>Acanthus mollis</i>
prince of wales juniper	<i>Juniperus horizontalis 'Prince of Wales'</i>
purple coneflower	<i>Echinacea purpurea</i>
roseum elegans rhododendron	<i>Rhododendron catawbiense 'Roseum elegans'</i>
spurge	<i>Euphorbia</i> spp.
stonecrop	<i>Sedum</i> spp.
summer phlox	<i>Phlox paniculata</i>
white festival baby's breath	<i>Gypsophila paniculata</i>
wine periwinkle	<i>Vinca minor 'Atropurpurea'</i>

**Snapshot 2.5 TG may be used on the following established plant species (Note: Limitations on recommended treatment methods):**



18/30

**Trees**

**Recommended Treatment Method:**

**C = Container Grown**

**F = Field Grown**

**Scientific Name**

**Common Name**

Scientific Name	Common Name	Treatment Method
<i>Abies balsamea</i>	fir, balsam	C, F
<i>Abies concolor</i>	fir, white	F
<i>Abutilon hybridum</i>	albus-flowering maple	C, F
	luteus-flowering maple	C, F
	roseus-flowering maple	C, F
	tangerine-flowering maple	C, F
	vesuvius red-flowering maple	F
<i>Acer ginnala</i>	flame maple	F
<i>Acer rubrum</i>	red maple	F
	red sunset maple	F
<i>Acer saccharinum</i>	silver maple	C, F
<i>Alsophila australis</i>	Australian tree fern	C, F
<i>Areacastrum romanzoffianum</i>	queen palm	C, F
<i>Betula nigra</i>	birch, river	C, F
<i>Betula papyrifera</i>	paper birch	F
<i>Brachychiton populneus</i>	bottle tree	C, F
<i>Bucida buceras</i>	black olive	F
<i>Ceratonia siliqua</i>	carob	F
<i>Cercis canadensis</i>	redbud	C, F
<i>Chamaecyparis obtusa</i>	filicoides-fernspray cypress	F
<i>Chamaecyparis obtusa</i>	gracilis-slender hinoki cypress	F
<i>Chamaecyparis pisifera</i>	sawara-false cypress	F
	squarrosa-moss cypress	F
<i>Chamaedorea cataractarum</i>	cat palm	F
	palm	C, F
<i>Chamaedorea costaricana</i>	palm	C, F
<i>Chamaedorea elegans</i>	parlor palm	C, F
<i>Cornus florida</i>	cloud nine dogwood	C, F
	dogwood, flowering	C, F
<i>Crataegus viridis</i>	green hawthorn	F
<i>Cryptomeria japonica</i>	cryptomeria, Japanese	C, F
<i>Cupaniopsis anacardioides</i>	carrot wood	F
<i>Cupressocyparis x 'Emerald Isle'</i>	emerald island leyland cypress	C, F
<i>Cupressus arizonica</i>	Arizona cypress	C, F
<i>Cupressus glabra</i>	Arizona cypress	F
<i>Cupressus sempervirens</i>	Italian cypress	C, F
<i>Dicksonia antarctica</i>	Tasmanian tree fern	C, F
<i>Elaeagnus angustifolia</i>	Russian olive	C, F
<i>Elaeagnus x 'Gilt edge'</i>	gilt edge silverberry	C, F
<i>Eucalyptus camaldulensis</i>	red gum eucalyptus	F
<i>Eucalyptus cinerea</i>	eucalyptus, mealy	F
	silver dollar eucalyptus	F
<i>Eucalyptus microtheca</i>	coolibah tree	C, F
<i>Eucalyptus sideroxylon</i>	eucalyptus, red ironbark	F
<i>Ficus benjamina</i>	figus	C, F
	mini ficus	C, F
<i>Fraxinus udhei</i>	shamel ash	C, F
<i>Ginkgo biloba</i>	ginkgo (maidenhair tree)	F
<i>Gleditsia triacanthos</i>	honey locust	F
	shademaster honey locust	F



<i>Acacia abyssinica</i>	glossy abelia	C, F	
<i>Acacia redolens</i>	abyssinica acacia	C, F	
<i>Acacia stenophylla</i>	acacia, prostrate	C, F	
<i>Acalypha wilkesiana</i>	shoestring acacia	C, F	
<i>Acer ginnala</i>	copper leaf	C, F	
<i>Acer palmatum</i>	amur maple	C, F	
	coral bark Japanese maple	C, F	
	dwarf Japanese maple	C, F	
<i>Agapanthus africanus</i>	queen anne lily of the Nile	C, F	
<i>Agave americana</i>	century plant	F	
<i>Anisodonteia hypomandarum</i>	cape mallow	C, F	
<i>Aptenia cordifolia</i>	red apple aptenia	C, F	
<i>Ardisia japonica</i>	chirimen marlberry	C, F	
<i>Astilbe arendsii</i>	false spiraea	C, F	
<i>Astilbe chinensis</i>	astilbe/false spirea	C, F	
<i>Athyrium nipponicum</i>	Japanese painted fern	C, F	
<i>Baccharis pilularis</i>	coyotebush	F	
<i>Berberis gladywensii</i>	william penn barberry	C, F	
<i>Berberis mentorensis</i>	mentor barberry	C, F	
<i>Berberis thunbergii</i>	aurea-golden Japanese barberry	F	
	crimson pygmy barberry	C, F	
	rose glow barberry	C, F	
	atropurea-redleaf Japanese barberry	F	
	cherry bomb barberry	C, F	
<i>Bougainvillea spp.</i>	barbara karst	C, F	
	California gold	C, F	
	pink pixie	C, F	
	scarlet o'hara	C, F	
	temple fire	C, F	
	Texas dawn	C, F	
<i>Buxus x 'Green velvet'</i>	green velvet boxwood	C, F	
<i>Buxus microphylla japonica</i>	boxwood, Japanese	C, F	
<i>Buxus microphylla Koreana</i>	Korean boxwood	F	
<i>Buxus sempervirens</i>	boxwood, common	C, F	
<i>Callistemon citrinus</i>	bottlebrush, lemon	F	
<i>Callistemon viminalis</i>	weeping bottlebrush	C, F	
<i>Calluna vulgaris</i>	spring torch Scotch heather	C, F	
<i>Camellia japonica</i>	camellia	C, F	
<i>Caryopteris x clandonen</i>	dark knight bluebeard	C, F	
<i>Cassia artemisioides</i>	cassia, feathery	C, F	
<i>Ceanothus spp.</i>	wild lilac	F	
<i>Cephalotaxus drupacae</i>	plum yew	C, F	
<i>Cerastium tomentosum</i>	snow-in-summer	C, F	
<i>Chamaecyparis obtusa</i>	kosteri cypress	C, F	
	nana-dwarf hinoki cypress	C, F	
	torulosa cypress	C, F	
<i>Chamaecyparis pisifera</i>	baileyi-dogwood	F	
	flaviramea-dogwood	F	
	squarrosa minima cypress	C, F	
<i>Chamaecyparis pisifera spp.</i>	filifera-thread cypress	C, F	
<i>Chrysalidocarpus lutescens</i>	areca palm	F	
<i>Clethra alnifolia</i>	summersweet	C, F	
<i>Cleyera japonica</i>	cleyera, Japanese	C, F	
<i>Coleonema pulchrum</i>	pink breath of heaven	C, F	
<i>Convolvulus cneorum</i>	bush morning glory	C, F	
<i>Cornus alba</i>	sibirica-Siberian dogwood	C, F	

2/30

<i>Cornus stolonifera</i>	baileyi-red-osier dogwood	F	
	flaviramea-yellowtwig dogwood	F	
<i>Cotinus coggygria</i>	royal purple smoke tree	C, F	
<i>Cotinus dammeri</i>	coral beauty smoke tree	C, F	
	eichholz smoke tree	C, F	
<i>Cotoneaster adpressus</i>	praecox-early cotoneaster	C, F	
<i>Cotoneaster apiculatus</i>	cotoneaster, cranberry	C, F	
<i>Cotoneaster congestus</i>	cotoneaster, Pyrenees	C, F	
<i>Cotoneaster dammeri</i>	cotoneaster, bearberry	C, F	
<i>Cotoneaster himalayan</i>	Himalayan cotoneaster	C, F	
<i>Cotoneaster horizontalis</i>	cotoneaster, rock	C, F	
<i>Cotoneaster opiculata</i>	cotoneaster	C, F	
<i>Cycas revoluta</i>	sago palm	C, F	
<i>Cytisus praecox</i>	hollandia-warminster broom	C, F	
<i>Cytisus scoparius</i>	lena-Scotch broom	C, F	
<i>Cytisus spp.</i>	holandia-Scotch broom	F	
<i>Daphne odora</i>	fragrant daphne	C, F	
<i>Deutzia crenata</i>	nakiana-dwarf deutzia	C, F	
<i>Deutzia gracilis</i>	slender gracilis	C, F	
<i>Dodonea viscosa</i>	hopseed bush	F	
<i>Elaeagnus pungens</i>	fruitland silver berry	C, F	
<i>Erica cinerea</i>	purple bell heather	C, F	
<i>Erica vagans</i>	cornish heather	C, F	
<i>Erica x darleyensa</i>	Mediterranean pink heather	C, F	
<i>Eugenia myrtifolia</i>	dwarf brush cherry	C, F	
<i>Euonymus x 'Aureo variegatus'</i>	gold spot euonymus	C, F	
<i>Euonymus x 'Chollipo'</i>	chollipo euonymus	C, F	
<i>Euonymus fortunei</i>	canadale gold euonymus	C, F	
	emerald'n gold euonymus	F	
	sunspot euonymus	C, F	
<i>Euonymus japonica</i>	silver king euonymus	F	
	variegated evergreen euonymus	C, F	
<i>Euonymus kiatschovica</i>	spreading euonymus	C, F	
<i>Euonymus vegetus</i>	bigleaf wintercreeper	C, F	
<i>Euryops pectinatus</i>	dwarf euryops	C, F	
<i>Fatsyhedera japonica</i>		C, F	
<i>Fatsia japonica</i>	Japanese aralia	C, F	
<i>Felicia ameloides</i>	blue marguerite	C, F	
<i>Forsythia intermedia</i>	forsythia, border	C, F	
<i>Forsythia x 'Spring glory'</i>	spring glory forsythia	C, F	
<i>Gardenia jasminoides</i>	august beauty gardenia	C, F	
	gardenia	C, F	
	radican gardenia	C, F	.....
<i>Gaultheria shallon</i>	salal/lemon leaf	C, F	.....
<i>Gelsemium sempervirens</i>	Carolina jessamine	C, F	.....
<i>Genista pilosa</i>	woadwaxen	C, F	.....
<i>Hibiscus rosa-sinensis</i>	ross estey-hibiscus	C, F	.....
<i>Hibiscus syriacus</i>	rose of sharon, red bird	C, F	.....
	rose of sharon, red heart	F	.....
	rose of sharon, woodbridge	C, F	.....
	rose of sharon, aphrodite		.....
	rose of sharon, helene		.....
<i>Ilex aquifolium</i>	Balkans holly	F	.....
	gold coast holly	F	.....
<i>Ilex aquipernyi</i>	san jose holly	C, F	.....





	goldfinger potentilla	C, F	
	red ace potentilla	C, F	
	sunset potentilla	C, F	
	tangerine potentilla	C, F	
<i>Potentilla verna</i>	spring cinquefoil	C, F	
<i>Prunus gladulosa</i>	dwarf pink flowering almond	C, F	
<i>Pyracantha fortuneana</i>	lolendei monrovia pyracantha	C, F	
	monon pyracantha	F	
	red elf hybrid pyracantha	F	
	rutgers hybrid pyracantha	C, F	
	santa cruz pyracantha	C, F	
	victory pyracantha	F	
<i>Rhaphiolepis indica</i>	charisma-monruce raphiolepis	C, F	
	enchantress-moness raphiolepis	F	
	raphiolepis (India hawthorn)	C, F	
	springtime-monme raphiolepis	F	
<i>Rhaphiolepis ovata</i>	roundleaf raphiolepis	C, F	
<i>Rhododendron calendulaceum</i>	cannon's double azalea	C, F	
	flame azalea	F	
	golden flare azalea	C, F	
	klondike azalea	C, F	
<i>Rhododendron campylocarpum</i>	butterfly rhododendron	F	
<i>Rhododendron carolinianum x daurium</i>	PJM rhododendron	C, F	
<i>Rhododendron catawbiense</i>	catawba album rhododendron	C, F	
	catawba rhododendron	C, F	
	lord roberts rhododendron	C, F	
	rocket rhododendron	C, F	
<i>Rhododendron caucasicum x ponticum</i>	cunningham white rhododendron	C, F	
<i>Rhododendron exbury</i>	cannon's double azalea	C, F	
	golden flare azalea	C, F	
	klondike azalea	C, F	
<i>Rhododendron forrestii repens</i>	gomer waterer rhododendron	C, F	
<i>Rhododendron forrestii x griersonianum</i>	elizabeth rhododendron	C, F	
<i>Rhododendron griffithianum</i>	jean marie rhododendron	C, F	
<i>Rhododendron hybrid spp.</i>	America rhododendron	C, F	
	English roseum rhododendron	F	
	nova zembla rhododendron	C, F	
	scintillation rhododendron	C, F	
<i>Rhododendron impeditum</i>	rhododendron	C, F	
<i>Rhododendron indica</i>	formosa azalea	C, F	
<i>Rhododendron indica</i>	waucabusa azalea	C, F	
<i>Rhododendron kaempferi</i>	blue danube azalea	C, F	
<i>Rhododendron kerume</i>	coral bells azalea	C, F	
	hino crimson azalea	C, F	
	hino pink azalea	C, F	
	mildred azalea	C, F	
	snow azalea	C, F	
<i>Rhododendron maximum</i>	rhodie max (rosebay)	C, F	
<i>Rhododendron mucronulatum</i>	rhododendron	F	
<i>Rhododendron obtusum</i>	hino-crimson azalea	C, F	
<i>Rhododendron ponticum</i>	chionoides rhododendron	C, F	
<i>Rhododendron ponticum</i>	daphnoides rhododendron	C, F	
<i>Rhododendron x 'purple gem'</i>	purple gem rhododendron	C, F	
<i>Rhododendron racemosum</i>	dwarf scarlet wonder rhododendron	C, F	
	tribly rhododendron	C, F	
	unique rhododendron	C, F	



<i>Viburnum setigerum</i>	tea viburnum	F
<i>Viburnum tinus compactum</i>	spring bouquet viburnum	F
<i>Viburnum trilobum</i>	cranberry bush	C, F
<i>Viburnum trilobum compactum</i>	dwarf cranberry bush	C, F
<i>Viburnum x pragense</i>	viburnum	C, F
<i>Weigela florida</i>	bristol ruby weigela	C, F
	java red weigela	C, F
	minuet weigela	C, F
<i>Xylosma congestum</i>	xylosma	F
<i>Yucca filamentosa</i>	yucca	C, F


**Groundcovers/Perennials**

**Recommended Treatment Method:**

**C = Container Grown**

**F = Field Grown**

Scientific Name	Common Name	
<i>Achillea millefolium</i>	common yarrow	C, F
<i>Agapanthus africanus</i>	lily of the Nile	C, F
<i>Agapanthus "Peter Pan"</i>		C, F
<i>Alstroemeria aurea</i>	Peruvian lily	C, F
<i>Ammophila breviligulata</i>	beechgrass	C, F
<i>Antirrhinum majus</i>	snapdragon	C, F
<i>Arctotheca calendula</i>	cape weed	F
<i>Argyranthemum frutescens</i>	Paris daisy	C, F
<i>Artemisia schmidtiana</i>	angels' hair	C, F
<i>Asparagus retrofractus</i>	fern	C, F
<i>Asteriscus maritimus</i>	gold coin daisy	C, F
<i>Astilbe Deutschland</i>	deutschland astilbe	C, F
<i>Asparagus retrofractus</i>		C, F
<i>Asparagus variegata</i>	tree fern	C, F
<i>Aster novae-angliae</i>	New England aster	C, F
<i>Aster novi-belgii</i>	New York aster	C, F
<i>Begonia cordifolia</i>	heartleaf begonia	C, F
<i>Begonia semperflorens</i>	white ambassador begonia	C, F
<i>Bidens ferulifolia</i>	peter's gold bidens	C, F
<i>Brachycome x 'New amethyst'</i>	swan river daisy new amethyst	C, F
<i>Callistephus chinensis</i>	China aster	C, F
<i>Carex spp.</i>	sedge	C, F
<i>Carpobrotus edulis</i>	ice plant, largeleaf	F
<i>Catharanthus roseus</i>	Madagascar periwinkle	C, F
<i>Cerastium tomentosum</i>	snow in the summer	C, F
<i>Ceratostigma plumbaginoides</i>	dwarf plumbago	C, F
<i>Chrysanthemum morifolium</i>	florist's chrysanthemum	C, F
<i>Chrysanthemum sp.</i>	chrysanthemum species	C, F
<i>Clematis integrifolia caerulea</i>	blue bell clematis	C, F
<i>Clivia miniata</i>	kafir lily	C, F
<i>Coreopsis verticillata</i>	coreopsis, threadleaf	C, F
<i>Cortaderia selloana</i>	pampas grass	C, F
<i>Cuphea hyssopifolia</i>	false or Mexican heather	C, F
<i>Cyperus albostratus</i>	dwarf umbrella grass	C, F
<i>Dahlia x 'Royal dahlietta pink'</i>	dwarf dahlia wendy pink	C, F
<i>Delosperma alba</i>	white iceplant	F
<i>Descampsia caespitosa</i>	descampsia	C, F
<i>Dianthus gratianopolitanus</i>	crimson treasure cheddar pink	C, F
<i>Dietes vegeta</i>	fortnight lily	C, F
<i>Drosanthemum floribundum</i>	trailing rosea iceplant	F





<i>Osteospermum fruticosum</i>	mondo grass	C, F
<i>Pachysandra terminalis</i>	freeway daisy	C, F
<i>Parthenocissus quinquefolia</i>	Japanese spurge	C, F
<i>Pelargonium x hortorum</i>	Virginia creeper	C, F
<i>Pelargonium peltatum</i>	zonal geranium	C, F
<i>Pennisetum alopecuroides</i>	ivy geranium	C, F
<i>Pennisetum setaceum</i>	fountain grass	C, F
<i>Pentas lanceolata</i>	chrimson fountaingrass	C, F
<i>Penstemon x 'Apple blossom'</i>	star cluster	C, F
<i>Penstemon gentianoides</i>	apple blossom penstemon	C, F
<i>Perovskia atriplicifolia</i>	hartwig penstemon	C, F
<i>Petunia-hybrids</i>	Russian sage	C, F
<i>Phalaris arundinacea picta</i>	garden petunias	C, F
<i>Ratibida columnifera</i>	ribbon grass	C, F
<i>Rudbeckia fulgida</i>	Mexican hat	C, F
<i>Rudbeckia hirta</i>	blackeyed susan	C, F
<i>Ruellia brittoniana</i>	blackeyed susan	C, F
<i>Salvia grahamii</i>	dwarf katie ruellia	C, F
<i>Salvia leucantha</i>	graham's sage	C, F
<i>Sedum x 'Autumn joy'</i>	Mexican bush sage	C, F
<i>Sedum x 'Vera jameson'</i>	autumn joy stonecrop	C, F
<i>Targetes patula 'Little hero'</i>	vera jameson stonecrop	C, F
<i>Trachelospermum asiaticum</i>	little hero marigold	C, F
<i>Tulbaghia violacea</i>	asian jasmine	C, F
<i>Verbena peruviana</i>	society garlic	C, F
<i>Vinca major</i>	st. paul verbena	C, F
<i>Vinca minor</i>	periwinkle, bigleaf	C, F
<i>Vinca spp.</i>	periwinkle, dwarf	F
<i>Zinnia elegans</i>	periwinkle	F
	dwarf zinnia	C, F

**Non-Bearing Fruit and Nut Trees**

**Non-bearing Vineyards<sup>1</sup>**

**Recommended Treatment Method:**  
**C = Container Grown**  
**F = Field Grown**

**Common Name**

almond	F	
apple	F	
apricot	F	
avocado	F	
blackberry	F	
blueberry	F	cccc
boysenberry	F	cccc
cherry, sour	F	cccc
cherry, sweet	F	cccc
currant	F	
dewberry	F	cccc
elderberry	F	cccc
fig	F	cccc
filbert	F	cccc
gooseberry	F	cccc
grape, American	F	cccc
grape, European	F	cccc
grapefruit	F	cccc

kiwi	F
lemon	F
loganberry	F
macadamia nut	F
nectarine	F
olive	F
orange	F
peach	F
pear	F
pecan	F
pistachio	F
plum	F
pomegranate	F
prune	F
raspberry	F
walnut, black	F
walnut, English	F

<sup>1</sup>Non-bearing fruit and nut trees and non-bearing vineyards are defined as plants which will not bear fruit for at least one year after treatment.

### Ornamental Bulbs

Do not use Snapshot 2.5 TG in bulb production fields. For use in landscape settings only.

Snapshot 2.5 TG may be applied for control of susceptible annual weeds in ornamental bulbs, e.g., bulbous iris, daffodil (narcissus), hyacinth and tulip. Apply Snapshot 2.5 TG to the soil surface 2 to 4 weeks after planting, but prior to the emergence of annual weeds. Snapshot 2.5 TG may also be applied following bulb emergence but prior to flowering and after flowering. For fall planted bulbs, apply Snapshot 2.5 TG again in late winter or early spring to weed-free soil surfaces.

#### Special Use Precautions:

Do not apply to tulip plants that have emerged to a height greater than 3/4 inch.

Do not apply to gladiolus prior to emergence or if corms are less than one inch in diameter.

Do not apply to bulbs while they are flowering.

### Shadehouse Areas

Snapshot 2.5 TG may be applied in open shadehouse-type structures where the natural flow of air is unimpeded. Do not apply in enclosed greenhouses or in enclosed shadehouse-type structures. Do not apply within three weeks prior to enclosing greenhouse or poly-type structures.

### Non-Cropland

Snapshot 2.5 TG is recommended as a preemergence treatment for control of certain broadleaf weeds and annual grasses on non-cropland areas such as industrial sites, utility substations, highway guardrails, sign posts, delineators, etc. Apply Snapshot 2.5 TG anytime prior to germination of target weeds. Areas to be treated should be free of established weeds or existing weeds should be controlled with postemergence herbicides.

Refer to the General Information section of this label for General Use Precautions and information on Application, Application Rates, and Weeds Controlled.

---

### Terms and Conditions of Use

---

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid.

Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

**Warranty Disclaimer**

Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Dow AgroSciences MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

**Inherent Risks of Use**

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. All such risks shall be assumed by buyer.

**Limitation of Remedies**

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Dow AgroSciences' election, one of the following:

- 1. Refund of purchase price paid by buyer or user for product bought, or
- 2. Replacement of amount of product used.

Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing. In no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

®Trademark of Dow AgroSciences LLC  
EPA accepted 03/20/02

